

Disorder and Localization in the Lowest Landau Level

Z. Gedik and M. Bayindir

Department of Physics, Bilkent University, Bilkent 06533 Ankara, Turkey

(March 7, 2009)

We study the localization property of a two-dimensional noninteracting electron gas in the presence of randomly distributed short-range scatterers. We evaluate the participation number of the eigenstates obtained by exact diagonalization technique. At low impurity concentrations we obtain self-averaged values showing that all states, except those exactly at the Landau level, are localized with finite localization length. We conclude that there is no universal localization exponent and at least at low impurity concentrations localization length does not diverge.

PACS numbers: 71.70.Di, 73.40.Hm, 71.23.-k, 72.15.Rn, 71.70.Di, 73.40.Hm

There has been a long lasting interest in understanding the localization problem in two-dimensional (2D) systems. According to scaling theory of localization [1–4], all states in a 2D system are localized if a disordered potential is present. However, in the presence of a strong perpendicular magnetic field, where the time reversal symmetry is broken, extended states appear in the center of impurity-broadened Landau bands [5,6]. If the scattering between Landau levels can be neglected, these extended states exist only at a single energy [7]. The width of the quantized plateaus of the integer quantum Hall effect (QHE) depends on the ratio of number of localized to extended states [8].

In analogy with the quantum critical phenomena and other localization transitions, it has been proposed that localization length $\xi(E)$ diverges as E approaches the critical energy E_c , which is equal to Landau level energy, so that

$$\xi(E) \propto |E - E_c|^{-\nu} \quad (1)$$

where ν is the localization critical exponent [9]. After the initial calculations of Aoki and Ando [10–12], several groups attempted to determine this critical exponent [13–27]. Experimental results [28–32] are generally in good agreement with the calculated values. Various techniques have allowed the computation of the exponent ν , and they strongly suggest a universal value close to $\nu = 7/3$ for the lowest Landau level (LLL). However, in spite of a great deal of experimental evidence and numerical simulations in its favor, there is no rigorous derivation of power law divergence in the localization length. Furthermore, even if the power law divergence is true, it is not clear whether the localization critical exponent is universal, independent of impurity concentration or parameters of the disordered potential [33].

Recently, we developed a method for a particle in the LLL moving in an arbitrary potential [34]. In this study we apply the method, which is basically an exact diagonalization technique, to a potential formed by randomly distributed short-range scatterers. We concentrate on

low impurity concentrations where it is difficult to perform calculations by other methods due to the presence of zero eigenvalues associated with the extended states at the band center. At low enough concentrations, we obtain self-averaged values where energy spectrum or localization property of eigenstates do not change with increasing system size. Contrary to the widely accepted view, localization length does not diverge at low impurity concentrations but instead the maximum localization length grows exponentially with impurity density. Extrapolation to less pure systems suggests that localization length can become as large as the sizes of the samples used in QHE experiments which explains the observed divergence in measurements.

The Hamiltonian for a particle of mass m and charge q , moving in 2D in the presence of magnetic field $\mathbf{B} = \nabla \times \mathbf{A}$ perpendicular to the plane and potential V , is given by $H = H_0 + V$ where

$$H_0 = \frac{1}{2m}(\mathbf{p} - \frac{q}{c}\mathbf{A})^2 \quad . \quad (2)$$

Using the symmetric gauge $\mathbf{A} = \frac{1}{2}\mathbf{B} \times \mathbf{r}$ and complex coordinates $z = X + iY = \sqrt{qB/2\hbar c}(x + iy)$ where $\mathbf{r} = (x, y)$, the unperturbed Hamiltonian can be written as $H_0 = \hbar\omega(a^\dagger a + 1/2)$ where $a^\dagger = -\partial/\partial z + z^*/2$. Since $[a, a^\dagger] = 1$, the energy eigenvalues are given by $E_n = \hbar\omega(n + 1/2)$ where $\omega = qB/2mc$ (q is assumed to be positive) and $n = 0, 1, 2, \dots$. When the magnetic field is very high the particle is confined into the LLL. This is a good approximation as long as the potential is small in comparison to Landau level splitting $\hbar\omega$. We are going to measure energies from the LLL so that $E_n = 0$.

Now, let us consider the potential

$$V(z, z^*) = V_0 \sum_i \delta(z - z_i) \delta(z^* - z_i^*) \quad (3)$$

where z_i denotes the position of the i th impurity in complex coordinates defined above. According to our method [34], to find the nonzero eigenvalues, the matrix to be diagonalized is

$$\langle i|\tilde{V}|j\rangle = \frac{V_0}{\pi} \exp(z_i z_j^* - |z_i|^2/2 - |z_j|^2/2) \quad . \quad (4)$$

Once $\tilde{V}$ is diagonalized, the eigenfunctions $\psi(z, z^*)$ of V can be constructed from $\tilde{\psi}_i$

$$\psi(z, z^*) = \sqrt{\frac{V_0}{\pi^2 E}} \sum_i \exp(z z_i^* - |z|^2/2 - |z_i|^2/2) \tilde{\psi}_i \quad . \quad (5)$$

We distinguish between the extended and the localized states via participation number P , which is the inverse of the mean fourth power of the amplitude [35,36]. Therefore, given a wave function ψ , P is defined as

$$P = \frac{[\int |\psi(\mathbf{r})|^2 d\mathbf{r}]^2}{\int |\psi(\mathbf{r})|^4 d\mathbf{r}}. \quad (6)$$

The participation number is a convenient quantity for distinguishing between localized and extended states since it takes a nonvanishing value for the former and becomes infinite for the latter. If a state is localized within a d -dimensional volume of average diameter D , P behaves as D^d irrespectively of the system size. For a plane wave it depends on the system size as L as L^d . In general, extended states lead to an effective dimensionality d^* , smaller than the real dimensionality d , which means that the states are not space-filling [37].

For $\tilde{\psi}$, participation number reduces to $\tilde{P} = 1/\sum_i |\tilde{\psi}_i|^4$ provided that $\tilde{\psi}$ is normalized. For a state localized on single impurity $\tilde{P} = 1$, while for uniform distribution over N_i impurities $\tilde{P} = N_i$. Therefore, we can interpret $\tilde{P}$ as a measure of number of scatterers on which $\tilde{\psi}$ takes nonzero value. We note that, as can be seen from Eqn. 5, ψ and $\tilde{\psi}$ have the same localization behavior, i.e. they are both extended or localized. Although we can evaluate corresponding P , we prefer to use $\tilde{P}$ to distinguish between localized and extended states. In this way we get rid of four-fold sums to be performed for evaluation of P in terms of $\tilde{\psi}_i$. We note that for $f < 1$, where f is the number of impurities per flux quantum, there are $N_i(1/f - 1)$ extended states at the center of the Landau band. However, the behavior given by Eqn. 1 has been proposed for states with $E \neq E_c$ and our method filters the states with $E = E_c$.

In Fig. 1, we plot participation number $\tilde{P}$ as a function of energy measured in units of V_0/π . We keep the concentration of impurities the same ($f = 2/3$) and evaluate the energy eigenvalues and the eigenfunctions for different distributions of the scatterers. For this example, $N_i = 1250$ and the number of configurations is 16 so that there are 2×10^4 points in the graph. Figure 1 shows that participation number is direct measure of localization length. $\tilde{P} = 1$ states, which are localized on a single impurity, occur mainly at $E = 1$ as we expect. The inset shows the details of the region around $E = 1$.




FIG. 1. Participation number $\tilde{P}$ as a function of energy (measured in units of V_0/π) for $f = 2/3$, i.e. two impurities per three flux quanta. There are $N_i = 1250$ impurities and 16 different distributions are used. The inset shows the details of the region around $E = 1$.

Localization around a purely repulsive scattering center is due to confinement into the LLL. All other states involve at least two scatterers so that $\tilde{P} \geq 2$. We use $\tilde{P}$ to decide whether a state is extended or not as follows. Let us consider N_i impurities distributed in a square so that their concentration, i.e. number of impurities per flux quantum, is f . If $\tilde{P} > \sqrt{N_i}$, then $\tilde{\psi}$ is nonvanishing on approximately $\sqrt{N_i}$ sites which means that it may extend from one side of the square to its opposite. Therefore, we assume that $\tilde{\psi}$ is an extended state. On the other hand, if $\tilde{P} < \sqrt{N_i}$, then $\tilde{\psi}$ has no chance to be extended. In this way we obtain the number of extended states N_e , for a given system composed of N_i impurities. Although $\tilde{P} > \sqrt{N_i}$, corresponding state can still be localized. Therefore, N_e is only an upper bound for the number of extended states. Our definition of extendedness become exact if the sites at which $\tilde{\psi}$ is nonvanishing form straight lines at least in one direction.

Figure 2 shows variation of N_e with N_i for different impurity concentrations. Each point is obtained in such a way that number of different configurations times N_i is 10^5 . Straight lines indicate that the two numbers are related by $N_e \propto N_i^y$ and the inset shows variation of y with f . We note that y depends upon impurity concentration. If we assume that the maximum localization length diverges as energy E approaches the critical value E_c (see Eqn. 1), then it is easy to show that $\nu = 1/2(1 - y)$. The widely accepted value $\nu = 7/3$ corresponds to $y = 11/14$ which we obtain at $f = 0.97$. At lower f values, ν becomes lower. However, we must be careful in comparing our results with the values in the literature since most, in fact to our knowledge all, of the existing calculations have been performed at impurity concentrations higher




FIG. 2. Number of extended states N_e for a given number of impurities N_i . Linear fitting gives a relation $N_e \propto N_i^y$. Each point is obtained in such a way that the number of different configurations times N_i is 10^5 . The inset shows the variation of y with impurity concentration f .

than unity. In addition to this, our definition of extend- edness is somewhat arbitrary. In general, the number of extended states is less than N_e . Finally, as we are going to see from the next figure, divergence of the localization length can merely be an artifact of finite size calculations.

As we go to more dilute systems and perform calcu- lations in large enough systems a remarkable change oc- curs. As shown in Fig. 3, N_e no more increases with N_i but instead decreases and vanishes at the end. For $N_i > N_i^*$ there are no extended states. We define N_i^* as the number of impurities for which $N_e = 1$. If we increase N_i further, we do not get any extended states. This re- sult is independent of the arbitrariness of our method of distinguishing between extended and localized states.

At this stage, it is not possible to say whether there is no localization length divergence for larger concen- trations but Fig. 4 gives some idea about the system sizes to be used if the same result holds in this denser region. In Fig. 4, we plot N_i^* as a function of f . We observe that N_i^* increases exponentially which suggests that for large enough sizes it may be possible to observe the same be- havior even for higher concentrations. If this is the case, then Fig. 2 is only an artifact of finiteness of the systems that we used in our calculations.

We also performed calculations for the case with equal number of attractive and repulsive point scatterers. We observe that there is no qualitative change in our results as can be seen from the figure. Beyond a certain size all states are localized and localization length does not diverge.

It is known that for very clean samples QHE is not observed [30]. Assuming that each impurity atom corre- sponds to a single δ -function potential, we evaluate the




FIG. 3. Number of extended states N_e vs number of im- purities N_i for $f = 2/3$. After $N_i = N_i^* = 1598$ no extended states appear.

concentration f but the resulting values (0.01-0.1) are too small. However, it should be noted that in our model scatterers are short-ranged while in the experiment op- posite is true, i.e. disorder potential is smoothly vary- ing in comparison to the magnetic length. In that case we can use a correlated disordered potential where sev- eral δ -functions are used to model the extended potential created by a single impurity atom. Another point that should be taken into account before quantitative com- parison is that at low impurity concentrations electron- electron interactions become important.




FIG. 4. Variation of N_i^* with f . The upper line is for pure repulsive case while the lower is obtained for the case with equal number of attractive and repulsive scattering centers.

In conclusion, at low impurity concentrations we ob- tain self-averaged values showing that all states, except those exactly at the Landau level, are localized with finite localization length. We conclude that the localization

exponent is not universal and at least at low impurity concentrations localization length does not diverge. Our results suggest that the same behavior can be observed for higher concentrations. If this is the case then there is no universal localization exponent, and in contrast to results of previous theoretical, numerical, and experimental studies, localization length does not diverge. On the other hand, if for higher concentrations there is a divergence, the transition between the two regimes is a very interesting problem to investigate.

ACKNOWLEDGMENTS

This work was supported by the Scientific and Techni- cal Research Council of Turkey (TUBITAK) under grant No. TBAG-1736. We gratefully acknowledge useful dis- cussions with Prof. I.O. Kulik.

-
- [1] D. J. Thouless, Phys. Rep. **13**, 93 (1974).
 - [2] F. Wegner, Z. Phys. B **25**, 327 (1976); *ibid* **35**, 207 (1979).
 - [3] E. Abrahams, P.W. Anderson, D.C. Licciardello, and T.V. Ramakrishnan, Phys. Rev. Lett. **42**, 673 (1979).
 - [4] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, Rev. Mod. Phys. **69**, 315 (1997).
 - [5] H. Aoki and T. Ando, Solid State Commun. **38**, 1079 (1981).
 - [6] Y. Ono, J. Phys. Soc. Jpn. **51**, 2055 (1982).
 - [7] J.T. Chalker, J. Phys. C **20**, L493 (1987).
 - [8] K. von Klitzing, G. Dorda, and M. Pepper, Phys. Rev. Lett. **45**, 494 (1980).
 - [9] S.A. Trugman, Phys. Rev. B **27**, 7539 (1983).
 - [10] T. Ando, J. Phys. Soc. Jpn. **52**, 1740 (1983).
 - [11] H. Aoki and T. Ando, Phys. Rev. Lett. **54**, 831 (1985).
 - [12] T. Ando and H. Aoki, J. Phys. Soc. Jpn. **54**, 2238 (1985).
 - [13] S. Hikami, Prog. Theor. Phys. **76**, 1210 (1986).
 - [14] J.T. Chalker and P.D. Coddington, J. Phys. C **21**, 2665 (1988).
 - [15] A.M.M. Pruisken, Phys. Rev. Lett. **61**, 1297 (1988).
 - [16] G.V. Milnikov and I.M. Sokolov, JETP Lett. **48**, 536 (1988).
 - [17] B. Huckestein and B. Kramer, Phys. Rev. Lett. **64**, 1437 (1990).
 - [18] B. Mielke, Europhys. Lett. **13**, 453 (1990).
 - [19] Y. Huo and R.N. Bhatt, Phys. Rev. Lett. **68**, 1375 (1992).
 - [20] U. Fastenrath, M. Janssen, and W. Pook, Physica A **191**, 401 (1992).
 - [21] D.H. Lee, Z. Wang, and S. Kivelson, Phys. Rev. Lett. **70**, 4130 (1993).
 - [22] H.L. Zhao and S. Feng, Phys. Rev. Lett. **70**, 4134 (1993).
 - [23] D. Liu and S. Das Sarma, Phys. Rev. B **49** 2677 (1994).
 - [24] T. Ando, Phys. Rev. B **49** 4679 (1994).
 - [25] B. Huckestein, Rev. Mod. Phys. **67**, 357 (1995).
 - [26] H. Matsuoka, Phys. Rev. B **55** R7327 (1997).
 - [27] I. Varga *et al.*, cond-mat/9710036.
 - [28] H.P. Wei, D. C. Tsui, M. A. Paalanen, and A. M. M. Pruisken, Phys. Rev. Lett. **61**, 1294 (1988).
 - [29] J. Wakabayashi, M. Yamane, and S. Kawaji, J. Phys. Soc. Jpn. **58**, 1903 (1989).
 - [30] S. Koch, R.J. Haug, K. von Klitzing, and K. Ploog, Phys. Rev. B **43**, 6828 (1991); *ibid* **46**, 1596 (1992). Phys. Rev. Lett. **67**, 883 (1991).
 - [31] H.P. Wei, L.W. Engel, and D.C. Tsui, Phys. Rev. B **50**, 14609 (1994).
 - [32] M. Furlan, Phys. Rev. B **57**, 14818 (1998).
 - [33] S. Das Sarma, in *Perspectives in Quantum Hall Effects*, edited by S.D. Sarma and A. Pinczuk, (Wiley, New York, 1997).
 - [34] Z. Gedik and M. Bayindir, Phys. Rev. B **56**, 12088 (1997).
 - [35] R.J. Bell and P. Dean, Discuss. Faraday Soc. **50**, 55 (1970).
 - [36] F. Wegner, Z. Phys. B **36**, 209 (1980).
 - [37] M. Schreiber, Phys. Rev. B **31**, 6146 (1985).